MULTI-SYMPTOM MAPAP COLD FORMULA- acetaminophen, dextromethorphan hbr, phenylephrine hcl tablet Major Pharmaceuticals

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Major 44-470C-CMS

Active ingredients (in each caplet)

Acetaminophen 325 mg Dextromethorphan HBr 10 mg Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer Cough suppressant Nasal decongestant

Uses

- temporarily relieves these common cold and flu symptoms:
 - minor aches and pains
 - nasal congestion
 - headache
 - cough
 - sinus congestion and pressure
 - sore throat
- helps clear nasal passages
- promotes nasal and sinus drainage
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- blisters
- rash

skin reddening

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease) or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- thyroid disease
- high blood pressure
- difficulty in urination due to enlargement of the prostate gland
- liver disease
- heart disease
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- diabetes
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- new symptoms occur
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- cough comes back or occurs with rash or headache that lasts. These could be signs
 of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed
- adults and children 12 years and over
 - take 2 caplets every 4 hours
 - swallow whole do not crush, chew, or dissolve
 - do not take more than 10 caplets in 24 hours
- children under 12 years: ask a doctor

Other information

- TAMPER EVIDENT: DO NOT USE IF OUTER PACKAGE IS OPENED OR BLISTER IS TORN OR BROKEN
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

corn starch, croscarmellose sodium, crospovidone, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, silicon dioxide, stearic acid, sucralose, talc, titanium dioxide

Questions or comments? (800) 616-2471

Principal Display Panel

MAJOR®

NDC 0904-5786-24 Compare to active ingredients in Tylenol® COLD MAX*

MULTI-SYMPTOM Mapap® COLD FORMULA CAPLETS

ACETAMINOPHEN

DEXTROMETHORPHAN HBr PHENYLEPHRINE HCI Pain Reliever / Fever Reducer Cough Suppressant • Nasal Decongestant

For Relief of: Fever, Headache, Sore Throat, Nasal Congestion, Coughing DAYTIME • NON-DROWSY ASPIRIN FREE

24 CAPLETS with Cool Blast Flavor

Actual Size

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Tylenol $^{\rm (B)}$ COLD MAX.

50844 REV0718F47008

Distributed by MAJOR® PHARMACEUTICALS

17177 N Laurel Park Drive, Suite 233

Livonia, MI 48152 USA M-17 Rev. 08/18

Re-order No. 006363

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

Nasal decongestant Phenylephine HCl 5 mg ■ µejbe cjest usesj bseædee Dextromethorphan HBr 10 mg....Cough suppressant a sinus congestion and pressure sore throat ostaminophen 325 mg ... Pain reliever/fever reducer headache moiteegn∞ lasen ■ ■ condp (in each caplet) and flu symptoms: ■ minor aches and pains Active ingredients temporarily relieves these common cold Purpose sasn

Drug Facts (continued)

COMPLETE PRO DUCT INFORMATION REPOUTER PACKAGE FOR

Drug Facts

TAMPER EVIDENT: DO NOT USE IF PACKAGE IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING

*This product is not manufactured or distributed by Johnson & Johnson Corporation, owner of the registered trademark Tylenol® COLD MAX. 50844 REV0718F47008

Rev. 08/18

Distributed by MAJOR® PHARMACEUTICALS 17177 N Laurel Park Drive, Suite 233 Livonia, MI 48152 USA M-17 Rev. 08/ Re-order No. 006363

MAJOR[®]

NDC 0904-5786-24 Compare to active ingredients in Tylenol® COLD MAX*

MULTI-SYMPTOM CAPLETS

ACETAMINOPHEN

DEXTROMETHORPHAN HBr PHENYLEPHRINE HCI

Pain Reliever / Fever Reducer Cough Suppressant • Nasal Decongestant

For Relief of: Fever, Headache, Sore Throat, Nasal Congestion, Coughing

DAYTIME • NON-DROWSY

ASPIRIN FREE

24 CAPLETS with Cool **Blast Flavor**

Actual Size

■ store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F) OR BLISTER IS TORN OR BROKEN DO NOT USE IF OUTER PACKAGE IS OPENED

Other information **TAMPER** EVIDENT:

Drug Facts (continued)

alcohol, povidone, silicon dioxide, stearic acid, sodium, crospovidone, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl Inactive ingredients corn starch, croscamellose

■ see end flap for expiration date and lot number

Drug Facts (continued)

B-1212-470C-08-CMS REV0718F47008

■ children under 12 years: ask a doctor

■ do not take more than 10 caplets in 24 hours ■ swallow whole - do not œush, chew, or dissolve

■ take 2 caplets every 4 hours

■ adults and children 12 years and over ■ do not take more than directed

Directions

children even if you do not notice any signs or medical attention is critical for adults as well as for

Control Center (1-800-222-1222) right away. Prompt overdose, get medical help or contact a Poison Keep out of reach of children. In case of accidental professional before use.

If pregnant or breast-feeding, ask a health

■ thyroid disease
■ high blood pressure Ask a doctor before use if you have

product or any of its ingredients ■ if you have ever had an allergic reaction to this

or pharmacist before taking this product. prescription drug contains an MAOI, ask a doctor the MAOI drug. If you do not know if your Parkinson's disease), or for 2 weeks after stopping depression, psychiatric or emotional conditions, or oxidase inhibitor (MANI) (certain drugs for ■ if you are now taking a prescription monoamine

doctor or pharmacist. sure whether a drug contains acetaminophen, ask a

■ with any other drug containing acetaminophen esu fon o O

doctor promptly.

(prescription or nonprescription). If you are not




sucralose, talc, titanium dioxide Drug Facts (continued)

Questions or comments? (800) 616-2471

_ot no. & Exp. Date No print area

WZ



taking the blood thinning drug wartamn. When using this product do not exceed recommended dosage.

Drug Facts (continued)

help right away. Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a


■ blisters ■ rash ■ skin reddening If a skin reaction occurs, stop use and seek medical

this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- Warnings
 Liver waming: The product contains acetaminophen.
 Severe liver damage may occur if you take
 more than 4,000 mg of acetaminophen in 24 hours
 with other drugs containing acetaminophen

 3 or more alcoholic drinks every day while using
 - promotes nasal and sinus drainage
 temporarily reduces fever

Drug Facts (continued)



Major 44-470C

MULTI-SYMPTOM MAPAP COLD FORMULA

acetaminophen, dextromethorphan hbr, phenylephrine hcl tablet

Product Information				
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0904-5786	
Route of Administration	ORAL			

Active Ingredient/Active Moiety				
Ingredient Name	Basis of Strength	Strength		
ACETAMINOPHEN (UNII: 36209ITL9D) (ACETAMINOPHEN - UNII:36209ITL9D)	ACETAMINOPHEN	325 mg		
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg		
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg		

Inactive Ingredients	
Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE, UNSPECIFIED (UNII: FZ 989GH94E)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
CROSPOVIDONE, UNSPECIFIED (UNII: 2S7830E561)	
POLYVINYL ALCOHOL, UNSPECIFIED (UNII: 532B59J990)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics					
Color	white	Score	no score		
Shape	OVAL	Size	17mm		
Flavor	MINT	Imprint Code	44;470		
Contains					

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:0904- 5786-24	2 in 1 CARTON	07/15/2005		
1		12 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	07/15/2005	

Labeler - Major Pharmaceuticals (191427277)

Establishment			
Name	Address	ID/FEI	Business Operations
LNK International, Inc.		832867894	manufacture(0904-5786)

Establishment				
Name	Address	ID/FEI	Business Operations	
LNK International, Inc.		832867837	pack(0904-5786)	

Revised: 3/2021 Major Pharmaceuticals